



LABORATORY OF FUNDAMENTAL AND APPLIED PHYSICS
CADI AYYAD UNIVERSITY, SAFI, MOROCCO



PROBING FOR A 95 GEV HIGGS BOSON WITHIN A 2-HIGGS DOUBLET MODEL

MOHAMMED BOUKIDI

IN COLLABORATION WITH: A. BELYAEV, R. BENBRIK, M. CHAKRABORTI, S. MORETTI AND S. SEMLALI

[arXiv:2306.09029 \[hep-ph\]](https://arxiv.org/abs/2306.09029)

THE FIRST EDITION OF THE AFRICAN CONFERENCE ON HIGH ENERGY PHYSICS

October 24, 2023

OUTLINE

1 INTRODUCTION

2 GENERAL TWO HIGGS DOUBLET MODEL

- 2HDM PARAMETRIZATION
- YUKAWA COUPLINGS
- ALIGNMENT LIMIT


3 NUMERICAL RESULTS

- RESULTS

4 CONCLUSION

INTRODUCTION

- ◆ Higgs properties measurements at run 1 and run 2 are in a good agreement with the SM
- ◆ Perhaps other scalars are not yet discovered
- ◆ Two Higgs Doublet Model (2HDM)
 - ◆ Minimal extension to the SM
 - ◆ Rich collider phenomenology
 - ◆ LHC benchmark mode
 - ◆ Benchmarks for light/heavy charged Higgs
 - ◆ Benchmarks for light /heavy neutral Higgses




C. Patrignani et al., Particle Physics Group, Chin. Phys. C, 40 100001

MOTIVATION

★ a 2.9σ local excess at ~ 95 GeV in $\gamma\gamma$ channel


$$\begin{aligned}\mu_{\gamma\gamma} &= \frac{\sigma(gg \rightarrow h)}{\sigma_{SM}(gg \rightarrow h)} \times \frac{\mathcal{BR}(h \rightarrow \gamma\gamma)}{\mathcal{BR}_{SM}(h \rightarrow \gamma\gamma)} \\ &= |c_{htt}|^2 \times \frac{\mathcal{BR}(h \rightarrow \gamma\gamma)}{\mathcal{BR}_{SM}(h \rightarrow \gamma\gamma)} \\ &= 0.33^{+0.19}_{-0.12}.\end{aligned}$$



[CMS-PAS-HIG-20-002]

★ a 2.6σ local excess at ~ 95 GeV in $\tau\tau$ channel

$$\begin{aligned}\mu_{\tau\tau} &= \frac{\sigma(gg \rightarrow h)}{\sigma_{SM}(gg \rightarrow h)} \times \frac{\mathcal{BR}(h \rightarrow \tau\tau)}{\mathcal{BR}_{SM}(h \rightarrow \tau\tau)} \\ &= |c_{htt}|^2 \times \frac{\mathcal{BR}(h \rightarrow \tau\tau)}{\mathcal{BR}_{SM}(h \rightarrow \tau\tau)} \\ &= 1.2 \pm 0.5.\end{aligned}$$



[CMS-HIG-21-001]

MOTIVATION


LHC Seminar

Measurement of Higgs boson production and search for new resonances in final states with photons and Z bosons, with the ATLAS detector

by Chiara Arcangeletti (INFN e Laboratori Nazionali di Frascati (IT))

Tuesday Jun 6, 2023, 10:00 AM → 11:00 AM Africa/Casablanca

500/1-001 - Main Auditorium (CERN)



$$\mu_{\gamma\gamma}^{ATLAS} = 0.18 \pm 0.10 \text{ (1.7}\sigma\text{)}$$

$$\mu_{\gamma\gamma}^{ATLAS+CMS} = 0.24^{+0.09}_{-0.08} \text{ (3.1}\sigma\text{)}$$

[T. Biekotter, S. Heinemeyer, G. Weiglein 2306.03889]

2HDM PARAMETRIZATION

The most general scalar potential of the 2HDM :

$$\begin{aligned}
 V(\Phi_1\Phi_2) = & \mathbf{m}_{11}^2 \Phi_1^\dagger \Phi_1 + \mathbf{m}_{22}^2 \Phi_2^\dagger \Phi_2 - \left[\mathbf{m}_{12}^2 \Phi_1^\dagger \Phi_2 + \text{h.c.} \right] \\
 & + \frac{\lambda_1}{2} \left(\Phi_1^\dagger \Phi_1 \right)^2 + \frac{\lambda_2}{2} \left(\Phi_2^\dagger \Phi_2 \right)^2 + \lambda_3 \left(\Phi_1^\dagger \Phi_1 \right) \left(\Phi_2^\dagger \Phi_2 \right) + \lambda_4 \left(\Phi_1^\dagger \Phi_2 \right) \left(\Phi_2^\dagger \Phi_1 \right) \\
 & + \left\{ \frac{\lambda_5}{2} \left(\Phi_1^\dagger \Phi_2 \right)^2 + \left[\lambda_6 \left(\Phi_1^\dagger \Phi_1 \right) + \lambda_7 \left(\phi_2^\dagger \Phi_2 \right) \right] \Phi_1^\dagger \Phi_2 + \text{h.c.} \right\}
 \end{aligned} \tag{1}$$

with :

$$\Phi_{1,2} = \begin{pmatrix} \phi_{1,2}^+ + i\varphi_{1,2}^+ \\ \frac{1}{\sqrt{2}} (v_{1,2} + \rho_{1,2} + i\eta_{1,2}) \end{pmatrix} \tag{2}$$

- ◆ The 10 independent parameters (\mathbf{m}_{11}^2 , \mathbf{m}_{22}^2 , \mathbf{m}_{12}^2 , $\lambda_{1,\dots,7}$) are assumed to be real.
- ◆ 2 minimization conditions and the combination $v_1^2 + v_2^2 \implies 7$ free parameters:

$$\mathbf{m}_h, \mathbf{m}_H, \mathbf{m}_A, \mathbf{m}_{H^\pm}, \alpha, \tan \beta = \frac{v_2}{v_1} \text{ and } \mathbf{m}_{1,2}^2.$$

YUKAWA COUPLINGS

- ◆ Tree-level FCNCs allowed \implies both doublets can couple to leptons and quarks.
- ◆ The associated model is called **2HDM type-III.**
- ◆ The Yukawa Lagrangian in terms of physical scalar masses:

$$\begin{aligned}
 -\mathcal{L}_Y^{III} = & \sum_{f=u,d,\ell} \frac{m_j^f}{v} \left[(\xi_h^f)_{ij} \bar{f}_{Li} f_{Rj} h + (\xi_H^f)_{ij} \bar{f}_{Li} f_{Rj} H - i(\xi_A^f)_{ij} \bar{f}_{Li} f_{Rj} A \right] \\
 & + \frac{\sqrt{2}}{v} \sum_{k=1}^3 \bar{u}_i \left[\left(m_i^u (\xi_A^{u*})_{ki} V_{kj} P_L + V_{ik} (\xi_A^d)_{kj} m_j^d P_R \right) \right] d_j H^+ \\
 & + \frac{\sqrt{2}}{v} \bar{\nu}_i (\xi_A^\ell)_{ij} m_j^\ell P_R \ell_j H^+ + h.c. ,
 \end{aligned} \tag{3}$$

- ◆ To get naturally small **FCNCs**, one can use the ansatz formulated by: $\tilde{Y}_{ij} \propto \sqrt{m_i m_j}/v \chi_{ij}$

ϕ	$(\xi_\phi^u)_{ij}$	$(\xi_\phi^d)_{ij}$	$(\xi_\phi^\ell)_{ij}$
h	$\frac{c_\alpha}{s_\beta} \delta_{ij} - \frac{c_{\beta-\alpha}}{\sqrt{2}s_\beta} \sqrt{\frac{m_i^u}{m_j^u}} \chi_{ij}^u$	$-\frac{s_\alpha}{c_\beta} \delta_{ij} + \frac{c_{\beta-\alpha}}{\sqrt{2}c_\beta} \sqrt{\frac{m_i^d}{m_j^d}} \chi_{ij}^d$	$-\frac{s_\alpha}{c_\beta} \delta_{ij} + \frac{c_{\beta-\alpha}}{\sqrt{2}c_\beta} \sqrt{\frac{m_i^\ell}{m_j^\ell}} \chi_{ij}^\ell$
H	$\frac{s_\alpha}{s_\beta} \delta_{ij} + \frac{s_{\beta-\alpha}}{\sqrt{2}s_\beta} \sqrt{\frac{m_i^u}{m_j^u}} \chi_{ij}^u$	$\frac{c_\alpha}{c_\beta} \delta_{ij} - \frac{s_{\beta-\alpha}}{\sqrt{2}c_\beta} \sqrt{\frac{m_i^d}{m_j^d}} \chi_{ij}^d$	$\frac{c_\alpha}{c_\beta} \delta_{ij} - \frac{s_{\beta-\alpha}}{\sqrt{2}c_\beta} \sqrt{\frac{m_i^\ell}{m_j^\ell}} \chi_{ij}^\ell$
A	$\frac{1}{t_\beta} \delta_{ij} - \frac{1}{\sqrt{2}s_\beta} \sqrt{\frac{m_i^u}{m_j^u}} \chi_{ij}^u$	$t_\beta \delta_{ij} - \frac{1}{\sqrt{2}c_\beta} \sqrt{\frac{m_i^d}{m_j^d}} \chi_{ij}^d$	$t_\beta \delta_{ij} - \frac{1}{\sqrt{2}c_\beta} \sqrt{\frac{m_i^\ell}{m_j^\ell}} \chi_{ij}^\ell$

ALIGNMENT LIMIT

In the Higgs-basis the alignment limit is most clearly exhibited :

$$H_1 = \begin{pmatrix} H_1^+ \\ H_1^0 \end{pmatrix} \equiv \Phi_1 \cos \beta + \Phi_2 \sin \beta, \quad H_2 = \begin{pmatrix} H_2^+ \\ H_2^0 \end{pmatrix} \equiv -\Phi_1 \sin \beta + \Phi_2 \cos \beta$$

$$H_1 = \begin{pmatrix} G^+ \\ (v + S_1 + iG^0) / \sqrt{2} \end{pmatrix}, \quad H_2 = \begin{pmatrix} H^+ \\ (S_2 + iS_3) / \sqrt{2} \end{pmatrix}$$

The 2 physical Higgs states h et H are as follows:

$$H = (\sqrt{2}\text{Re}H_1^0 - v)\cos(\beta - \alpha) + \sqrt{2}\text{Re}H_2^0 \sin(\beta - \alpha) \quad (4)$$

$$h = (\sqrt{2}\text{Re}H_1^0 - v)\sin(\beta - \alpha) + \sqrt{2}\text{Re}H_2^0 \cos(\beta - \alpha) \quad (5)$$

- ◆ $\cos(\beta - \alpha) \rightarrow 0, h \equiv H_{SM}$ (J. Bernon, J. F. Gunion, H. E. Haber, Y. Jiang and S. Kraml, Phys. Rev. D 92 (2015) no.7, 075004): Inverted hierarchy
- ◆ $\sin(\beta - \alpha) \rightarrow 0, H \equiv H_{SM}$ (J. Bernon, J. F. Gunion, H. E. Haber, Y. Jiang and S. Kraml, Phys. Rev. D 93 (2016) no.3, 035027): Standard hierarchy

CONSTRAINTS

Theoretical

- ★ **Unitarity** constraints require a variety of scattering process to be unitary: specifically, the tree-level 2-to-2 body scattering matrix involving scalar-scalar, gauge-gauge and/or scalar-gauge initial and/or final states must have eigenvalues e_i 's such that $|e_i| < 8\pi$.
- ★ **Perturbativity** constraints impose the following condition on the quartic couplings of the scalar potential: $|\lambda_i| < 8\pi$
- ★ **Vacuum stability** constraints require the potential be bounded from below and positive in any direction of the fields Φ_i , consequently, the parameter space must satisfy the following conditions:

$$\begin{aligned}\lambda_1 > 0, \quad \lambda_2 > 0, \quad \lambda_3 > -\sqrt{\lambda_1 \lambda_2}, \\ \lambda_3 + \lambda_4 - |\lambda_5| > -\sqrt{\lambda_1 \lambda_2}.\end{aligned}$$

2HDMC-1.8.0 (D. Eriksson, J. Rathsman and O. Stal [0902.0851])

Experimental


- ★ **EWPOs**, implemented through the EW oblique parameters S, T , we require $\Delta\chi^2(S, T) \leq 6.18$.
- ★ **SM-like Higgs boson discovery**: an agreement between selected points in parameter space and the current measurements of the properties of the discovered Higgs boson at 125 GeV is enforced by means of the publicly available code [HiggsSignals-3 via HiggsTools\[2210.09332\]](#) (P. Bechtle et al).
- ★ **Non-SM-like Higgs boson exclusions**: to check the parameter space points against the exclusion limits from null Higgs boson searches at LEP, Tevatron and, in particular, the LHC, we apply the public code [HiggsBounds-6 via HiggsTools](#) (P. Bechtle et al).
- ★ **B-physics observables** are tested against data by resorting to the public code [SuperIso_v4.1 \(F. Mahmoudi \[0808.3144\]\)](#), (mainly $B \rightarrow X_s \gamma$, $B_{s,d} \rightarrow \mu^+ \mu^-$ and $B_s \rightarrow \tau \nu$).

NUMERICAL RESULTS


2HDM TYPE-III INTERPRETATION

m_h	m_H	m_A	m_{H^\pm}	$s_{\beta-\alpha}$	$\tan \beta$	m_{12}^2	$\chi_{ij}^{f,\ell}$
[94; 97]	125.09	[80; 300]	[160; 300]	[-0.5; 0.5]	[1; 30]	$m_h^2 \tan \beta / (1 + \tan^2 \beta)$	[-3; 3]

$\Delta \chi^2_{125} \leq 5.99$
 ----- 1 σ C.L. for χ^2_{x+y}
 + $\mu_{\gamma\gamma, \tau\tau}^{\text{exp}}$
 ★ $\min(\chi^2_{\gamma\gamma + \tau\tau})$
 ☆ $\min(\chi^2_{125})$




2HDM TYPE-III INTERPRETATION



2HDM TYPE-III INTERPRETATION

$$\Delta\chi^2_{125} \leq 5.99$$

- 1 σ C.L. (CMS)
- 1 σ C.L. (HL-LHC)
- 2 σ C.L. (HL-LHC)
- 1 σ C.L. (ILC 500)
- 2 σ C.L. (ILC 500)
- ★ min($\chi^2_{\gamma\gamma+\tau\tau}$)
- ★ min(χ^2_{125})
- ◆ SM



CONCLUSION

- ◆ Two local excesses ($\gamma\gamma$: 2.9σ + $\tau\tau$: 2.6σ) at the same mass ≈ 95 GeV.
- ◆ The generic 2HDM type III can accommodate both excesses simultaneously.
- ◆ We explored how our model could be studied at upcoming colliders such as the HL-LHC and ILC, focusing on the h_{125} couplings.

Thanks!

BACK-UP

